

# **SIMVASTATIN** Brand names: Zocor®

## What is this medication used for?

Simvastatin is used to lower "bad" cholesterol and triglycerides in the blood. It lowers the risk of stroke, heart attack, and other heart complications in people with diabetes, coronary heart disease, or other risk factors.

## How should I take this medication?

Simvastatin is taken by mouth.

Simvastatin should be taken at bedtime or with the evening meal. If you take it more than once per day, take it with meals.

If you miss a dose, take it as soon as you remember. If it is almost time for your next dose, skip the missed dose and wait until your next regularly scheduled dose. Do not take extra medicine to make up for the missed dose.

Do not take any new medication (prescription or over-the-counter) without checking with your doctor or pharmacist first.

### What are the side effects of this medication?

- headache
- mild muscle pain
- mild joint pain
- constipation
- mild nausea
- stomach pain or indigestion
- mild skin rash
- sleep problems (insomnia)

Other side effects not listed may also occur. Check with your physician if you notice any other effects.

#### Contact your doctor immediately if you experience any of the following while taking this medicine:

- unexplained muscle pain, tenderness, or weakness
- fever, unusual tiredness, and dark colored urine
- pain or burning when you urinate
- swelling, weight gain, urinating less than usual or not at all
- nausea, upper stomach pain, itching, loss of appetite, clay-colored stools, jaundice (yellowing of the skin or eyes)

## **Special Instructions:**

Your blood will need to be tested often to make sure that simvastatin is not causing any harmful effects. Your liver function may also need to be tested. Visit your doctor regularly and keep your appointments for these tests.

You may need to take simvastatin on a long-term basis for the treatment of high cholesterol.

This information is only a summary. Please contact your healthcare professional for additional information about this medication.